

MITES (ARACHNIDA: ACARI) INHABITING COFFEE DOMATIA: A SHORT REVIEW AND RECENT FINDINGS FROM COSTA RICA

Fernando E. Vega^{1,*}, Ronald Ochoa², Carlos Astorga³ and David E. Walter⁴

1. Sustainable Perennial Crops Laboratory (* corresponding author, e-mail: Fernando.Vega@ars.usda.gov) and
2. Systematic Entomology Laboratory, United States Department of Agriculture, Agricultural Research Service, BARC-W, Beltsville, Maryland 20705, U.S.A (e-mail: Ron.Ochoa@ars.usda.gov); 3. Unidad de Recursos Fitogenéticos y Biotecnología, Departamento de Agricultura y Agroforestería, Centro Agronómico Tropical de Investigación y Enseñanza, CATIE 7170, Turrialba, Costa Rica; 4. Department of Biological Sciences, CW 405, Biological Sciences Centre, University of Alberta, Edmonton, Alberta, Canada T6G 2E9 (DEW@ualberta.ca).

ABSTRACT - Eight previously unreported domatia-inhabiting mites are reported from *Coffea arabica* L. and *C. eugenioides* S. Moore (Rubiaceae) accessions planted in Costa Rica. One of these, an *Asca* sp., was found to be carrying fungal spores on its cuticle. A review of the literature on mites in coffee domatia is presented.

Internat. J. Acarol. 33(4): 291-295.