

## TWO NEW SPECIES OF *TETRANYCHUS* (ACARI: TETRANYCHIDAE) INFESTING CORN IN INNER MONGOLIA, CHINA

Shôzô Ehara<sup>1</sup>, Tetsuo Gotoh<sup>2</sup> and Xiao-Yue Hong<sup>3</sup>

1. Hamasaka 2-15-7, Tottori 680-0001, Japan (e-mail: shozoehama@ncn-t.net); 2. Laboratory of Applied Entomology and Zoology, Faculty of Agriculture, Ibaraki University, Ami, Ibaraki 300-0393, Japan (e-mail: gotoh@mx.ibaraki.ac.jp); 3. Department of Entomology, Nanjing Agricultural University, Nanjing, Jiangsu 210095, China (e-mail: xyhong@njau.edu.cn).

**ABSTRACT** - *Tetranychus huhhotensis* n. sp. and *T. zae* n. sp. are described and illustrated from corn leaves in Inner Mongolia. *Tetranychus huhhotensis* n. sp., assigned to Flechtmann and Knihinicki's Group 4 of *Tetranychus*, can be separated from its allied species by having the aedeagal knob with dorsal surface distinctly emarginate centrally. *Tetranychus zae* n. sp. is distinctive in Group 9 of this genus in that the aedeagal knob is minute, with anterior projection practically absent and posterior projection conspicuous, acute. Summer females of the two new species are pale greenish-yellow, while their diapausing females are pale-orange.

Internat. J. Acarol. 34(4): 339-345.